Infusion Therapy - IV Remdesivir Referral Form

- Patients will receive treatment in our community nursing clinics, unless under exceptional circumstances.
- We process only completed referrals (legible, signed, dated). Fax to 613.745.6984 or 1.855.450.8569.

Name		[DOB					HCN /	' VC		
Address								Unit			
City		Postal Code									
Phone		P	Alt Pho	one							
Preferred language for service: EN □ FR □ Other □ (specify)											
Diagnosis											
Allergies											
If applicable	harge Date						tion Control Precautions are LET, AIRBORNE and CONTACT				
 ☐ Use alternate contact (instead of patient) for assessment, due to ☐ Preference ☐ Hearing ☐ Cognition ☐ Language ☐ Other (specify) 											
Alt Contact	Name		Rela	tionsh	nip to pt				Phone		
If any answers to the questions below are "No", we are unable to administer the first											
dose of IV Remdesivir in the community.										Yes	No
Has the prescriber confirmed the patient does not have any serious allergies / adverse											
reactions to the ordered medication or related drugs?											
Has the prescriber confirmed the patient does not have anaphylaxis to Remdesivir or anaphylaxis of unknown origin?											
Is the patient at least 18 years old?											
For six hours after receiving the first dose and should an adverse reaction occur, does the											
patient have access to a working telephone to call 911 or to a hospital within approximately											
30 minutes drive from medication administration address?											
To monitor the patient for adverse reactions for six hours after the medication is											
administered, the patient / SDM understands that a capable adult (18 years or older) should											
be present in the home or with the patient. 1) Patient qualifies for Remdesivir treatment, per Ontario Health and Ministry of Health quidelines.											
1) Patient qualifies for Remdesivir treatment, per Ontario Health and Ministry of Health guidelines as they do not require hospitalization; AND cannot take Paxlovid (nirmatrelvir and ritonavir), e.g.,											
	due to a drug interaction or contraindication.										
2) Dat	2) Date of COVID-19 symptom onset Date of positive test										
Leatment (Da aft	y 0 first day of symptoms and Day 1 first full day										
aft.	fter the day the symptoms started).										
3) L Remdesivir 200 mg IV on Day 1, 100 mg IV daily on days 2 and 3. All doses via peripheral IV.											
4) Is this a first dose? ☐ Yes ☐ No.											
If n	no, Dose 1 date & time Dose 2 date & time										
☐ I confirmed that the patient does not have any serious allergies or adverse reactions to the ordered or											
related medications.											
☐ I confirmed there are no contraindications to patient receiving IV Remdesivir in the community,											
including review of recent bloodwork (Cr, ALT, AST & eGFR within three months), hepatic and renal											
function, pregnancy/breastfeeding status.											
I explained the risks of having the first dose in the community to the patient / most responsible person and the patient / most responsible person has given verbal consent.											
Additional Information / Orders											
Physician/NP Name (please print)											
Signature		1			Dat	ا م					
If delegate						one nun	ıber				
responsible					MRP phone number for urgent situations						

Confidential when completed. If you received this form in error, please call us at 1.800.538.0520.