

☐ Kirkland Lake
53 Government Rd. W./
53, ch. Government O.
Kirkland Lake ON P2N 2E5

Tel/Tél : 705 567 2222
Fax/Télec : 705 567 9407

☐ North Bay
1164 Devonshire Ave./
1164, av. Devonshire
North Bay ON P1B 6X7

Tel/Tél : 705 476 2222
Fax/Télec : 705 474 0080

☐ Parry Sound
6 Albert St./
6, rue Albert
Parry Sound ON P2A 3A4

Tel/Tél : 1 800 440 6762
Fax/Télec : 1-855-773-4056

☐ Sault Ste. Marie
390 Bay St., Main Floor/
390, rue Bay, 1e étage
Sault Ste. Marie ON P6A 1X2

Tel/Tél : 705 949 1650
Fax/Télec : 705 949 1663

☐ Sudbury
40 Elm St., Suite 41-C/
40, rue Elm, bureau 41-C
Sudbury ON P3C 1S8

Tel/Tél : 705 522 3461
Fax/Télec : 705 522 3855

☐ Timmins
330 Second Ave., Suite 101/
330 av. Second, bureau 101
Timmins ON P4N 8A4

Tel/Tél : 705 267 7766
Fax/Télec : 705 267 7795

TTY / ATS 711 (ask operator for 1-888-533-2222 / veuillez demander le téléphoniste pour le 1-888-533-222)
Toll Free / Sans frais: 1-800-461-2919 or / ou 310-2222 no area code required / indicatif régional non requis.

***Hospital: Use hospital Ontario Health atHome fax number**

Negative Pressure Wound Therapy Referral Form

Name:		Health Card #:		Version Code:	
Address:				Postal Code:	
Date of Birth:		Phone:			
Gender: ☐ Male ☐ Female ☐ Non-binary ☐ Unknown Pronouns:					
Diagnosis:				Diabetic: ☐ Yes ☐ No	
Allergies: ☐ Yes ☐ No ☐ Unknown Specify:			Latex Allergy: ☐ Yes ☐ No ☐ Unknown		
WOUND TYPE					
The following conditions can be considered for the application of NPWT. Please indicate reason for referral.					
Acute Wound	☐ Surgical (dehiscid)	☐ Traumatic	☐ Abdominal	☐ Pilonidal cyst	☐ Partial thickness burn
Chronic Open Wound	☐ Diabetic ulcer (offloaded) ☐ Venous leg ulcer ☐ Stage 3 or 4 pressure injury (offloaded)				
Adjunct to Surgery	☐ Preparation of wound bed ☐ Incisional support ☐ Securing skin graft post-operatively				
Oncology Related	☐ Wound complicated by radiation ☐ Support wound healing prior to start of chemotherapy				
WOUND DESCRIPTION					
Location:		Length: cm x Width: cm x Depth: cm			
☐ Undermining Details if applicable:		☐ Tunneling Details if applicable:			
Note: NPWT will continue to be assessed in the community, and settings may be reviewed based on exudate and patient tolerance. Continuation of NPWT is dependent on wound healing goals being met. Maximum treatment time for NPWT is 8 weeks.					
NPWT TREATMENT ORDERS					
☐ ActiVAC (indicate pressure settings and dressing details below) Pressure (mmHg): _____ ☐ Continuous OR ☐ Intermittent Dressing (select one): Granufoam Black: ☐ Small (10cm x 7.5cm x 3.2cm) ☐ Medium (18cm x 12.5cm x 3.2cm) ☐ Large (26cm x 15cm x 3.2cm) ☐ X-Large (60cm x 30cm x 3.2cm) White Foam: ☐ Small (10cm x 7.5cm x 1cm) ☐ Large (10cm x 15cm x 1cm)			☐ PICO (single use, disposable) Pressure: ☐ 80 mmHg (non-adjustable) Dressing Size: ☐ 10cm x 20cm ☐ 10cm x 30cm ☐ 15cm x 15cm ☐ SNAP (single use, disposable) Pressure: ☐ 125 mmHg (non-adjustable) Dressing Size: ☐ 10cm x 10cm ☐ 15cm x 15cm V.A.C Peel and Place Small (up to 2cm in depth) Medium (up to 4cm in depth) Large (up to 6cm in depth)		

Name:		Health Card #:		Version Code:	
CONVENTIONAL DRESSING ORDERS					
Patients will be started on conventional dressings until NPWT can be initiated. Conventional orders also required in the case of service interruption.					
PRECAUTIONS AND CONTRAINDICATIONS					
The precautions and contraindications listed below have been reviewed, and it is determined that NPWT is appropriate to be used for patient ☐ YES ☐ NO (conventional dressings will be utilized until addressed)					
The following conditions are considered precautions in the use of NPWT: • Immunodeficiency (e.g. Leukemia, HIV); • Hematologic disorders; • Systemic or local signs of infection; • Uncontrolled diabetes; • Systemic steroids; • Receiving anticoagulant therapy; • The location of the wound will interfere with the therapy; • Nutritional impairment; • History of non-compliance; • Home environment not conducive to NPWT (i.e. cleanliness, animals etc.); or • Patient unable to adhere to minimum of 22 hours of therapy/day.			The following risk factors contraindicate the use of NPWT: • Inadequate wound visualization; • Untreated infection in the wound site; • Fistulas to body cavities or organs; • Presence of undebried necrotic tissue with eschar; • Untreated Osteomyelitis; • Malignancy or cancer in the wound margins; • Unresolved bleeding following debridement; or • Exposed vasculature, nerves or organ		
PRESCRIBER INFORMATION					
Name:		Phone:		Fax:	
Signature:		CPSO/CNO#:		Date:	
				After Hours Number:	