

Negative Pressure Wound Therapy Referral Form

[Clear Form](#)

Name:	Health Card Number:	Version Code:
Address:	Postal Code:	
Date of Birth:	Phone:	
Gender: ☐ Male ☐ Female ☐ Non-binary ☐ Unknown	Pronouns:	
Diagnosis:	Diabetic: ☐ Yes ☐ No	
Allergies: ☐ Yes ☐ No ☐ Unknown Specify:	Latex Allergy: ☐ Yes ☐ No ☐ Unknown	

WOUND TYPE

The following conditions can be considered for the application of NPWT. Please indicate reason for referral.

Acute Wound	☐ Surgical (dehiscid)	☐ Traumatic	☐ Abdominal	☐ Pilonidal cyst	☐ Partial thickness burn
Chronic Open Wound	☐ Diabetic ulcer (offloaded) ☐ Venous leg ulcer ☐ Stage 3 or 4 pressure injury (offloaded)				
Adjunct to Surgery	☐ Preparation of wound bed ☐ Incisional support ☐ Securing skin graft post-operatively				
Oncology Related	☐ Wound complicated by ☐ Support wound healing prior to starting chemotherapy				

WOUND DESCRIPTION

Location:	Length: cm x Width: cm x Depth: cm
☐ Undermining Details if applicable:	☐ Tunneling Details if applicable:

Note: NPWT will continue to be assessed in the community, and settings may be reviewed based on exudate and patient tolerance. Continuation of NPWT is dependent on wound healing goals being met. Maximum treatment time for NPWT is 8 weeks.

NPWT TREATMENT ORDERS

☐ ActiV.A.C. (indicate pressure settings and dressing details below) Pressure (mmHg): _____ ☐ Continuous OR ☐ Intermittent Dressing Type/Size (select one): Granufoam Black: ☐ Small (10cm x 7.5cm x 3.2cm) ☐ Medium (18cm x 12.5cm x 3.2cm) ☐ Large (26cm x 15cm x 3.2cm) ☐ X-Large (60cm x 30cm x 3.2cm) Simplace Ex: ☐ Small (7.7cm x 11.2cm x 1.75cm) ☐ Medium (14.7cm x 17.4cm x 1.75cm) White Foam: ☐ Small (10cm x 7.5cm x 1cm) ☐ Large (10cm x 15cm x 1cm)	☐ PICO (single use, disposable) Pressure: ☐ 80 mmHg (non-adjustable) Dressing Size (select one): ☐ 10cm x 20cm ☐ 15cm x 20cm ☐ 10cm x 30cm ☐ 15cm x 30cm ☐ 15cm x 15cm ☐ 20cm x 20cm ☐ SNAP (single use, disposable) Pressure: ☐ 125 mmHg (non-adjustable) Dressing Size (select one): ☐ 10cm x 10cm ☐ 15cm x 15cm
---	--

CONVENTIONAL DRESSING ORDERS

Patients will start on conventional dressings until NPWT can be initiated. Conventional orders are also required in the case of service interruption.

Patient Name:	Health Card No.:
---------------	------------------

PRECAUTIONS AND CONTRAINDICATIONS

The precautions and contraindications listed below have been reviewed, and it is determined that NPWT is appropriate to be used for the patient.

☐ **YES** ☐ **NO** (conventional dressings will be utilized until addressed)

The following conditions are considered precautions in the use of NPWT:

- Immunodeficiency (e.g. Leukemia, HIV);
- Hematologic disorders;
- Systemic or local signs of infection;
- Uncontrolled diabetes;
- Systemic steroids;
- Receiving anticoagulant therapy;
- The location of the wound will interfere with the therapy;
- Nutritional impairment;
- History of non-compliance;
- Home environment not conducive to NPWT (i.e. cleanliness, animals etc.); or
- Patient unable to adhere to minimum of 22 hours of therapy/day.

The following risk factors contraindicate the use of NPWT:

- Inadequate wound visualization;
- Untreated infection in the wound site;
- Fistulas to body cavities or organs;
- Presence of unbridged necrotic tissue with eschar;
- Untreated Osteomyelitis;
- Malignancy or cancer in the wound margins;
- Unresolved bleeding following debridement; or
- Exposed vasculature, nerves or organ

PRESCRIBER INFORMATION

Name:	Phone:	Fax:	After Hours Number:
Signature:	CPSO/CNO#:	Date:	